

DEPRESSION

Magnesium
Deficiency damages NMDA (N-methyl-D-aspartate) receptors in the brain, which regulate mood; Well-documented anti-depressant effects.^{1,2,3,4}

Zinc
Improves efficacy of antidepressant drugs; Particularly useful for treatment resistant patients; Regulates neurotransmitters.^{33,34,35,36}

Serine
Regulates brain chemistry; Involved in NMDA receptor function; Acts as a neurotransmitter; Low levels correlate with severity of depression.^{31,32}

Antioxidants
Oxidative stress in the brain alters neurotransmitter function; Antioxidants protect our brain, which is very sensitive to oxidation; Several antioxidants – Vitamins A, C and E, Lipoic Acid, CoQ10, Glutathione and Cysteine – play a key role in prevention and treatment of depression.^{28,29,30}

Biotin
Part of the B-vitamin complex, biotin deficiency has induced depression in animal and human studies.^{26,27}

Selenium
Integral part of regulatory proteins (selenoproteins) in the brain; Supplementation trials are promising; May alleviate postpartum depression.^{5,6}

Chromium
Elevates serotonin (feel-good neurotransmitter) levels in the brain; May be particularly effective on eating symptoms of depression such as carbohydrate craving and increased appetite, due to its effect on blood sugar regulation.^{37,38,39}

Folate
Building block for many “feel-good” neurotransmitters such as serotonin, dopamine and norepinephrine; Low folate causes poor response to antidepressant meds; The lower the folate, the more severe the depression.^{7,8,9,10}

Vitamin B12
Depression may be a manifestation of B12 deficiency; Repletion of B12 to adequate levels can improve treatment response; B12 deficiency common in psychiatric disorders.^{11,12,13}

Vitamin B6
Cofactor for serotonin and dopamine production (feel good chemicals); Studies indicate that low levels may predispose people to depression.^{14,15,16}

Vitamin B2
Low B2 has been implicated in depression due to its role in methylation reactions in the brain.^{17,18}

Vitamin D
Clinical trials suggest increasing blood levels of vitamin D, which is actually a hormone precursor, may improve symptoms of depression.^{19,20,21}

Inositol
Influences signaling pathways in the brain; Particularly effective in SSRI (selective serotonin reuptake inhibitor) sensitive disorders.^{24,25}

Carnitine
Increases serotonin and noradrenaline which lift mood; In trials, carnitine alleviates depression with few, if any, side effects.^{22,23}

REFERENCES

- ¹Nechifor M. Magnesium in major depression. *Magnes Res* 2009;22:163S-166S.
- ²Eby G, Eby K. Magnesium for treatment-resistant depression: a review and hypothesis. *Med Hypotheses* 2010;74:649-660.
- ³Szewczyk B, Poleszak E, Sowa-Kucma M et al. Antidepressant activity of zinc and magnesium in view of the current hypotheses of antidepressant action. *Pharmacol Rep* 2008;60:588-589.
- ⁴Eby G, Eby K. Rapid recovery from major depression using magnesium treatment. *Med Hypotheses* 2006;67:362-70.
- ⁵Mokhber N, Namjoo M, Tara F et al. Effect of supplementation with selenium on postpartum depression: a randomized double-blind placebo-controlled trial. *J Matern Fetal Neonatal Med* 2011;24:104-108.
- ⁶Benton D, Cook R. The impact of selenium supplementation on mood. *Biol Psychiatry* 1991;29:1092-1098.
- ⁷Coppen A, Bailey J. Enhancement of the antidepressant action of fluoxetine by folic acid: a randomized, placebo controlled trial. *J Affect Disord* 2000;60:121-130.
- ⁸Miller A. The methylation, neurotransmitter and antioxidant connections between folate and depression. *Altern Med Rev* 2008;13:216-226.
- ⁹Papakostas G, Petersen T, Mischoulon D et al. Serum folate, vitamin B12, and homocysteine in major depressive disorder, Part 1: predictors of clinical response in fluoxetine-resistant depression. *J Clin Psychiatry* 2004;65:1090-1095.
- ¹⁰Wesson V, Levitt A, Joffe R. Change in folate status with antidepressant treatment. *Psychiatry Res* 1994;53:313-322.
- ¹¹Kate N, Grover S, Agarwal M. Does B12 deficiency lead to a lack of treatment response to conventional antidepressants? *Psychiatry* 2010;74:42-44.
- ¹²Hintikka J, Tolmunen T, Tanskanen A et al. High vitamin B12 level and good treatment outcome may be associated in major depressive disorder. *BMC Psychiatry* 2003;3:17.
- ¹³Bhat A, Srinivasan K, Kurpad S et al. Psychiatric presentations of vitamin B12 deficiency. 2007;105:395-396.
- ¹⁴Merete C, Falcon L, Tucker K. Vitamin B6 is associated with depressive symptomatology in Massachusetts elders. *J Am Coll Nutr* 2008;27:421-427.
- ¹⁵Williams A, Cotter A, Sabina A et al. The role for vitamin B-6 as treatment for depression: a systematic review. *Fam Pract* 2005;22:532-537.
- ¹⁶Hvas A, Juul S, Bech P et al. Vitamin B6 level is associated with symptoms of depression. *Psychother Psychosom* 2004;73:340-343.
- ¹⁷Miyake Y, Sasaki S, Tanaka K et al. Dietary folate and vitamins B12, B6 and B2 intake and the risk of postpartum depression in Japan: the Osaka Maternal and Child Health Study. *J Affect Disord* 2006;96:133-138.
- ¹⁸Bell I, Morrow F, Read M et al. Low thyroxine levels in female psychiatric in patients with riboflavin deficiency: implications for folate-dependent methylation. *Acta Psychiatr Scand* 1992;85:360-363.
- ¹⁹Jorde R, Sneve M, Figenschau Y et al. Effects of vitamin D supplementation on symptoms of depression in overweight and obese subjects: randomized double blind trial. *J Intern Med* 2008;264:599-609.
- ²⁰Högberg G, Gustafsson SA, Hällström T et al. Depressed adolescents in a case-series were low in vitamin D and depression was ameliorated by vitamin D supplementation. *Acta Paediatr* 2012;101:779-783.
- ²¹Berk M, Sanders K, Pasco J et al. Vitamin D deficiency may play a role in depression. *Med Hypotheses* 2007;69:1316-1319.
- ²²Smeland O, Meisingset T, Borges K et al. Chronic acetyl-L-carnitine alters brain energy metabolism and increases noradrenaline and serotonin content in healthy mice. *Neurochem Int* 2012;61:100-107.
- ²³Garzya G, Corallo D, Fiore A et al. Evaluation of the effects of L-acetylcarnitine on senile patients suffering from depression. *Drugs Exp Clin Res* 1990;16:101-106.
- ²⁴Levine J, Barak Y, Gonzalves M et al. Double-blind, controlled trial of inositol treatment of depression. *Am J Psychiatry* 1995;152:792-794.
- ²⁵Brink C, Viljoen S, de Kock S et al. Effects of myo-inositol versus fluoxetine and imipramine pretreatments on serotonin 5HT2A and muscarinic acetylcholine receptors in human neuroblastoma cells. *Metab Brain Dis* 2004;19:51-70.
- ²⁶Levenson J. Biotin-responsive depression during hyperalimentionation. *J Parenter Enteral Nutr* 1983;7:181-183.
- ²⁷Osada K, Komai M et al. Experimental study of fatigue provoked by biotin deficiency in mice. *Int J Vitam Nutr Res* 2004;74:334-340.
- ²⁸Bodnar L, Wisner K. Nutrition and depression: Implications for improving mental health among child-bearing-aged women. *Biol Psychiatry* 2005;58:679-685.
- ²⁹Maes M, Mihaylova I, Kubera M et al. Lower plasma Coenzyme Q10 in depression: a marker for treatment resistance and chronic fatigue in depression and a risk factor to cardiovascular disorder in that illness. *Neuro Endocrinol Lett* 2009;30:462-469.
- ³⁰Tsuboi H, Shimoi K, Kinae N et al. Depressive symptoms are independently correlated with lipid peroxidation in a female population: comparison with vitamins and carotenoids. *J Psychosom Res* 2004;56:53-58.
- ³¹de Koning T, Klomp L. Serine-deficiency syndromes. *Curr Opin Neurol* 2004;17:197-204.
- ³²Mitani H, Shirayama Y, Yamada T et al. Correlation between plasma levels of glutamate, alanine and serine with severity of depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2006;30:1155-1158.
- ³³Cope E, Levenson C. Role of zinc in the development and treatment of mood disorders. *Curr Opin Clin Nutr Metab Care* 2010;13:685-689.
- ³⁴Nowak G, Siwek M, Dudek D et al. Effect of zinc supplementation on antidepressant therapy in unipolar depression: a preliminary placebo-controlled study. *Pol J Pharmacol* 2003;55:1143-1147.
- ³⁵Maes M, Vandoolaeghe E, Neels H et al. Lower serum zinc in major depression is a sensitive marker of treatment resistance and of the immune/inflammatory response in that illness. *Psychiatry* 1997;42:349-358.
- ³⁶Levenson C. Zinc, the new antidepressant? *Nutr Rev* 2006;64:39-42.
- ³⁷Attenburrow M, Odontiadis J, Murray B et al. Chromium treatment decreases the sensitivity of 5-HT2A receptors. *Psychopharmacology* 2002;159:432-436.
- ³⁸Docherty J, Sack D, Roffman M et al. A double-blind, placebo-controlled, exploratory trial of chromium picolinate in atypical depression: effect on carbohydrate craving. *J Psychiatr Pract* 2005;11:302-314.
- ³⁹Davidson J, Abraham K, Connor K et al. Effectiveness of chromium in atypical depression: a placebo-controlled trial. *Biol Psychiatry* 2003;53:261-264.

Additional references at <http://www.spectracell.com/online-library-mnt-depression-abstract/>